

THE INCLUSION RELATION BETWEEN SOBOLEV AND MODULATION SPACES

MASAHARU KOBAYASHI AND MITSURU SUGIMOTO

ABSTRACT. The inclusion relations between the L^p -Sobolev spaces and the modulation spaces is determined explicitly. As an application, mapping properties of unimodular Fourier multiplier $e^{i|D|^\alpha}$ between L^p -Sobolev spaces and modulation spaces are discussed.

1. INTRODUCTION

The modulation spaces $M_s^{p,q}$ are one of the function spaces introduced by Feichtinger [4] in 1980's to measure the decaying and regularity property of a function or distribution in a way different from L^p -Sobolev spaces L_s^p or Besov spaces $B_s^{p,q}$. The precise definitions of these function spaces will be given in Section 2, but the main idea of modulation spaces is to consider the space variable and the variable of its Fourier transform simultaneously, while they are treated independently in L^p -Sobolev spaces and Besov spaces.

Because of this special nature, modulation spaces are now considered to be suitable spaces in the analysis of pseudo-differential operators after a series of important works [3], [7], [8], [9], [17], [18] and so on. On the other hand, modulation spaces have also remarkable applications in the analysis of partial differential equations. For example, the Schrödinger and wave propagators, which are not bounded on neither L^p nor $B_s^{p,q}$, are bounded on $M_s^{p,q}$ ([2]). Modulation spaces are also used as a regularity class of initial data of the Cauchy problem for non-linear evolution equations, and in this way the existence of the solution is shown under very low regularity assumption for initial data (see [21], [22], [23]).

In the last several years, many basic properties of modulation spaces are established. In particular, the inclusion relation between Besov spaces and modulation spaces has been completely determined. Let us define the indices $\nu_1(p, q)$ and $\nu_2(p, q)$ for $1 \leq p, q \leq \infty$ in the following


Date: September 9, 2010.

way:

$$\nu_1(p, q) = \begin{cases} 0 & \text{if } (1/p, 1/q) \in I_1^* : \min(1/p, 1/p') \geq 1/q, \\ 1/p + 1/q - 1 & \text{if } (1/p, 1/q) \in I_2^* : \min(1/q, 1/2) \geq 1/p', \\ -1/p + 1/q & \text{if } (1/p, 1/q) \in I_3^* : \min(1/q, 1/2) \geq 1/p, \end{cases}$$

$$\nu_2(p, q) = \begin{cases} 0 & \text{if } (1/p, 1/q) \in I_1 : \max(1/p, 1/p') \leq 1/q, \\ 1/p + 1/q - 1 & \text{if } (1/p, 1/q) \in I_2 : \max(1/q, 1/2) \leq 1/p', \\ -1/p + 1/q & \text{if } (1/p, 1/q) \in I_3 : \max(1/q, 1/2) \leq 1/p, \end{cases}$$

where $1/p + 1/p' = 1 = 1/q + 1/q'$. We remark $\nu_2(p, q) = -\nu_1(p', q')$.



Then the following result is known:

THEOREM 1.1 (Sugimoto-Tomita [15], Toft [18]). *Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Then we have*

- (1) $B_s^{p,q}(\mathbf{R}^n) \hookrightarrow M^{p,q}(\mathbf{R}^n)$ if and only if $s \geq n\nu_1(p, q)$;
- (2) $M^{p,q}(\mathbf{R}^n) \hookrightarrow B_s^{p,q}(\mathbf{R}^n)$ if and only if $s \leq n\nu_2(p, q)$.

As for the inclusion relation between L^p -Sobolev spaces and modulation spaces, the following result (see also [19]) is immediately obtained from Theorem 1.1 if we notice the inclusion property $L_{s+\varepsilon}^p \hookrightarrow B_s^{p,q} \hookrightarrow L_{s-\varepsilon}^p$ for $\varepsilon > 0$ (see, [20, p.97]):

COROLLARY 1.2. *Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Then we have*

- (1) $L_s^p(\mathbf{R}^n) \hookrightarrow M^{p,q}(\mathbf{R}^n)$ if $s > n\nu_1(p, q)$. Conversely, if $L_s^p(\mathbf{R}^n) \hookrightarrow M^{p,q}(\mathbf{R}^n)$, then $s \geq n\nu_1(p, q)$;
- (2) $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$ if $s < n\nu_2(p, q)$. Conversely, if $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$, then $s \leq n\nu_2(p, q)$.

But in Corollary 1.2, there still remains a question whether the critical case $s = n\nu_1(p, q)$ or $s = n\nu_2(p, q)$ is sufficient or not for the inclusion. The objective of this paper is to answer this basic question and complete the picture of inclusion relations between the L^p -Sobolev spaces and the modulation spaces. The following theorems are our main results:

THEOREM 1.3. *Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Then $L_s^p(\mathbf{R}^n) \hookrightarrow M^{p,q}(\mathbf{R}^n)$ if and only if one of the following conditions is satisfied:*

- (1) $q \geq p > 1$ and $s \geq n\nu_1(p, q)$;
- (2) $p > q$ and $s > n\nu_1(p, q)$;
- (3) $p = 1, q = \infty$, and $s \geq n\nu_1(1, \infty)$;
- (4) $p = 1, q \neq \infty$ and $s > n\nu_1(1, q)$.

THEOREM 1.4. *Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Then $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$ if and only if one of the following conditions is satisfied:*

- (1) $q \leq p < \infty$ and $s \leq n\nu_2(p, q)$;
- (2) $p < q$ and $s < n\nu_2(p, q)$;
- (3) $p = \infty, q = 1$, and $s \leq n\nu_2(\infty, 1)$;
- (4) $p = \infty, q \neq 1$, and $s < n\nu_2(\infty, q)$.

It should be mentioned that Kobayashi-Miyachi-Tomita [10] determines the inclusion relation between modulation spaces $M_s^{p,q}$ and local Hardy spaces h^p for $0 < p \leq 1$. Our main results extend this result to the case $p > 1$ since we have $h^p = L^p$ then. As a matter of fact, the proof of Theorems 1.3 and 1.4 heavily depends on the results and arguments established in [10].

As an application of our main theorems, we also consider mapping properties of unimodular Fourier multiplier $e^{i|D|^\alpha}$, $\alpha \geq 0$, which is a generalization of wave ($\alpha = 1$) and Schrödinger ($\alpha = 2$) propagators. See Corollaries 5.2 and 5.4 in Section 5. As Theorem A and Theorem B there say, the operator $e^{i|D|^\alpha}$ ($0 \leq \alpha \leq 2$) is bounded on modulation spaces while not on L^p -Sobolev spaces. Theorems 1.3 and 1.4 help us to understand what happen if we consider the operator between L^p -Sobolev spaces and modulation spaces.

We explain the organization of this paper. After the next preliminary section devoted to the definitions and basic properties of function spaces treated in this paper, we give a proof of Theorem 1.4 in Sections 3 and 4. We remark that Theorem 1.3 is just the dual statement of Theorem 1.4. In Section 5, we consider mapping properties of unimodular Fourier multipliers between L^p -Sobolev spaces and modulation spaces, as well as those of invertible pseudo-differential operators.

2. PRELIMINARIES

2.1. Basic notation. The following notation will be used throughout this article. We write $\mathcal{S}(\mathbf{R}^n)$ to denote the Schwartz space of all complex-valued rapidly decreasing infinitely differentiable functions on \mathbf{R}^n and $\mathcal{S}'(\mathbf{R}^n)$ to denote the space of tempered distributions on \mathbf{R}^n , i.e., the topological dual of $\mathcal{S}(\mathbf{R}^n)$. The Fourier transform is defined by $\widehat{f}(\xi) = \int_{\mathbf{R}^n} f(x) e^{-ix \cdot \xi} dx$ and the inverse Fourier transform by $f^\vee(x) = (2\pi)^{-n} \widehat{f}(-x)$. We define

$$\|f\|_{L^p} = \left(\int_{\mathbf{R}^n} |f(x)|^p dx \right)^{1/p}$$

for $1 \leq p < \infty$ and $\|f\|_{L^\infty} = \text{ess. sup}_{x \in \mathbf{R}^n} |f(x)|$. We also define the L^p -Sobolev norm $\|\cdot\|_{L_s^p}$ by

$$\|f\|_{L_s^p} = \|(\langle \cdot \rangle^s \widehat{f}(\cdot))^\vee\|_{L^p} \quad \text{with} \quad \langle \cdot \rangle = (1 + |\cdot|^2)^{1/2}.$$

Let $(X, \|\cdot\|_X)$ and $(Y, \|\cdot\|_Y)$ be two Banach spaces, which include $\mathcal{S}(\mathbf{R}^n)$, respectively. We say that an operator T from X to Y is bounded if there exists a constant $C > 0$ such that $\|Tf\|_Y \leq C\|f\|_X$ for all $f \in \mathcal{S}(\mathbf{R}^n)$, and we set

$$\|T\|_{X \rightarrow Y} = \sup\{\|Tf\|_Y \mid f \in \mathcal{S}(\mathbf{R}^n), \|f\|_X = 1\}.$$

We use the notation $I \lesssim J$ if I is bounded by a constant times J , and we denote $I \approx J$ if $I \lesssim J$ and $J \lesssim I$.

2.2. Modulation spaces. We recall the modulation spaces. Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $\varphi \in \mathcal{S}(\mathbf{R}^n)$ be such that

$$(1) \quad \text{supp } \varphi \subset [-1, 1]^n \quad \text{and} \quad \sum_{k \in \mathbf{Z}^n} \varphi(\xi - k) = 1 \text{ for all } \xi \in \mathbf{R}^n.$$

Then the modulation space $M_s^{p,q}(\mathbf{R}^n)$ consists of all tempered distributions $f \in \mathcal{S}'(\mathbf{R}^n)$ such that the norm

$$\|f\|_{M_s^{p,q}} = \left(\sum_{k \in \mathbf{Z}^n} \langle k \rangle^{sq} \left(\int_{\mathbf{R}^n} |\varphi(D - k)f(x)|^p dx \right)^{q/p} \right)^{1/q}$$

is finite, with obvious modifications if p or $q = \infty$. Here we denote $\varphi(D - k)f(x) = (\varphi(\cdot - k)\widehat{f}(\cdot))^\vee(x)$.

We simply write $M^{p,q}(\mathbf{R}^n)$ instead of $M_0^{p,q}(\mathbf{R}^n)$ when $s = 0$. The space $M_s^{p,q}(\mathbf{R}^n)$ is a Banach space which is independent of the choice of $\varphi \in \mathcal{S}(\mathbf{R}^n)$ satisfying (1) ([4, Theorem 6.1]). If $1 \leq p, q < \infty$, then $\mathcal{S}(\mathbf{R}^n)$ is dense in $M_s^{p,q}(\mathbf{R}^n)$ ([4, Theorem 6.1]). If $1 \leq p_1 \leq p_2 \leq \infty$, $1 \leq q_1 \leq q_2 \leq \infty$ and $s_1 \geq s_2$ then $M_{s_1}^{p_1, q_1}(\mathbf{R}^n) \hookrightarrow M_{s_2}^{p_2, q_2}(\mathbf{R}^n)$ ([4, Proposition 6.5]). Let us define by $\mathcal{M}_s^{p,q}(\mathbf{R}^n)$ the completion of $\mathcal{S}(\mathbf{R}^n)$

under the norm $\|\cdot\|_{M_s^{p,q}}$. The dual of $\mathcal{M}_s^{p,q}(\mathbf{R}^n)$ can be identified with $\mathcal{M}_{-s}^{p',q'}(\mathbf{R}^n)$, where $1/p + 1/p' = 1 = 1/q + 1/q'$. If $1 \leq p, q < \infty$, then $\mathcal{M}_s^{p,q}(\mathbf{R}^n) = M_s^{p,q}(\mathbf{R}^n)$ ([1, Lemma 2.2]). Moreover, the complex interpolation theory for these spaces reads as follows: Let $0 < \theta < 1$ and $1 \leq p_1, p_2, q_1, q_2 \leq \infty, s_1, s_2 \in \mathbf{R}$. Set $1/p = (1-\theta)/p_1 + \theta/p_2$, $1/q = (1-\theta)/q_1 + \theta/q_2$ and $s = (1-\theta)s_1 + \theta s_2$, then $(\mathcal{M}_{s_1}^{p_1,q_1}, \mathcal{M}_{s_2}^{p_2,q_2})_{[\theta]} = \mathcal{M}_s^{p,q}$ ([4, Theorem 6.1], [21, Theorem 2.3]).

We recall the following lemmas.

LEMMA 2.1 ([4, Proposition 6.7]). *Let $1 \leq p \leq \infty$, $1/p + 1/p' = 1$ and $s \in \mathbf{R}$. Then*

$$M_s^{p,\min(p,p')}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n) \hookrightarrow M_s^{p,\max(p,p')}(\mathbf{R}^n).$$

Let $U_\lambda : f(x) \mapsto f(\lambda x)$ be the dilation operator. Then the following dilation property of $M^{p,q}$ is known.

LEMMA 2.2 ([15, Theorem 3.1]). *Let $1 \leq p, q \leq \infty$. We have, for $C_1, C_2 > 0$,*

$$\begin{aligned} \|U_\lambda f\|_{M^{p,q}} &\leq C_1 \lambda^{n\mu_1(p,q)} \|f\|_{M^{p,q}}, \quad \forall \lambda \geq 1, \forall f \in M^{p,q}(\mathbf{R}^n), \\ \|U_\lambda f\|_{M^{p,q}} &\geq C_2 \lambda^{n\mu_2(p,q)} \|f\|_{M^{p,q}}, \quad \forall \lambda \geq 1, \forall f \in M^{p,q}(\mathbf{R}^n), \end{aligned}$$

where

$$\begin{aligned} \mu_1(p, q) &= \begin{cases} -1/p & \text{if } (1/p, 1/q) \in I_1^* : \min(1/p, 1/p') \geq 1/q, \\ 1/q - 1 & \text{if } (1/p, 1/q) \in I_2^* : \min(1/q, 1/2) \geq 1/p', \\ -2/p + 1/q & \text{if } (1/p, 1/q) \in I_3^* : \min(1/q, 1/2) \geq 1/p, \end{cases} \\ \mu_2(p, q) &= \begin{cases} -1/p & \text{if } (1/p, 1/q) \in I_1 : \max(1/p, 1/p') \leq 1/q, \\ 1/q - 1 & \text{if } (1/p, 1/q) \in I_2 : \max(1/q, 1/2) \leq 1/p', \\ -2/p + 1/q & \text{if } (1/p, 1/q) \in I_3 : \max(1/q, 1/2) \leq 1/p. \end{cases} \end{aligned}$$

2.3. Besov spaces. We recall the Besov spaces. Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. Suppose that $\psi_0, \psi \in \mathcal{S}(\mathbf{R}^n)$ satisfy $\text{supp } \psi_0 \subset \{\xi \mid |\xi| \leq 2\}$, $\text{supp } \psi \subset \{\xi \mid 1/2 \leq |\xi| \leq 2\}$ and $\psi_0(\xi) + \sum_{j=1}^{\infty} \psi(\xi/2^j) = 1$ for all $\xi \in \mathbf{R}^n$. Set $\psi_j(\cdot) = \psi(\cdot/2^j)$ if $j \geq 1$. Then the Besov space $B_s^{p,q}(\mathbf{R}^n)$ consists of all $f \in \mathcal{S}'(\mathbf{R}^n)$ such that

$$\|f\|_{B_s^{p,q}} = \left(\sum_{j=0}^{\infty} 2^{jsq} \|(\widehat{f}(\cdot) \psi_j(\cdot))^\vee\|_{L^p}^q \right)^{1/q} < \infty,$$

with usual modification again if $q = \infty$.

The dual of $B_s^{p,q}(\mathbf{R}^n)$ can be identified with $B_{-s}^{p',q'}(\mathbf{R}^n)$, where $1/p + 1/p' = 1 = 1/q + 1/q'$.

2.4. Local Hardy spaces. We recall the local Hardy spaces. Let $0 < p < \infty$, and let $\Psi \in \mathcal{S}(\mathbf{R}^n)$ be such that $\int_{\mathbf{R}^n} \Psi(x) dx \neq 0$. Then the local Hardy space $h^p(\mathbf{R}^n)$ consists of all $f \in \mathcal{S}'(\mathbf{R}^n)$ such that

$$\|f\|_{h^p} = \left\| \sup_{0 < t < 1} |\Psi_t * f| \right\|_{L^p} < \infty,$$

where $\Psi_t(x) = t^{-n} \Psi(x/t)$. We remark that $h^1(\mathbf{R}^n) \hookrightarrow L^1(\mathbf{R}^n)$ ([5, Theorem 2]), $h^p(\mathbf{R}^n) = L^p(\mathbf{R}^n)$ if $1 < p < \infty$ ([5, p.30]), and the definition of $h^p(\mathbf{R}^n)$ is independent of the choice of $\Psi \in \mathcal{S}(\mathbf{R}^n)$ with $\int_{\mathbf{R}^n} \Psi(x) dx \neq 0$ ([5, Theorem 1]). The complex interpolation theory for these spaces reads as follows: Let $1 \leq p_1, p_2 < \infty$ and $0 < \theta < 1$. Set $1/p = (1 - \theta)/p_1 + \theta/p_2$, then $(h^{p_1}, h^{p_2})_{[\theta]} = h^p$ ([20, p.45]).

LEMMA 2.3 ([10]). *Let $1 \leq q \leq \infty$ and $s \in \mathbf{R}$. Then $h^1(\mathbf{R}^n) \hookrightarrow M_s^{1,q}(\mathbf{R}^n)$ if and only if $s \leq -n/q$. However, in the case $q \neq \infty$, $L^1(\mathbf{R}^n) \hookrightarrow M_s^{1,q}(\mathbf{R}^n)$ only if $s < -n/q$.*

3. SUFFICIENT CONDITIONS

We prove the *if* part of Theorem 1.4. First we remark the following fact:

LEMMA 3.1. *Let $1 < p \leq 2$, $p \leq q \leq p'$ and $s \leq -n(1/p + 1/q - 1)$. Then $L^p(\mathbf{R}^n) \hookrightarrow M_s^{p,q}(\mathbf{R}^n)$.*

PROOF OF LEMMA 3.1. We note that $L^2(\mathbf{R}^n) = M^{2,2}(\mathbf{R}^n)$ and, by Lemma 2.3,

$$h^1(\mathbf{R}^n) \hookrightarrow M_{-n/q}^{1,q}(\mathbf{R}^n)$$

for $1 \leq q \leq \infty$. The complex interpolation method yields

$$L^p(\mathbf{R}^n) \hookrightarrow M_{-n(1/p+1/q-1)}^{p,q}(\mathbf{R}^n),$$

which gives the desired result. \square

PROOF OF THEOREM 1.4 (“IF” PART). Suppose $q \leq p$ and $s \leq n\nu_2(p, q)$. If $q \leq \min(p, p')$, then $s \leq n\nu_2(p, q) = 0$ and we have

$$M^{p,q}(\mathbf{R}^n) \hookrightarrow M_s^{p,\min(p,p')}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$$

by Lemma 2.1. If $2 < p < \infty$ and $p' \leq q \leq p$, then $s \leq n\nu_2(p, q) = n(1/p + 1/q - 1)$ and we have $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$ again by the dual statement of Lemma 3.1. Thus we have the sufficiency of conditions (1) and (3). Conditions (2) and (4) are sufficient by Corollary 1.2. \square

4. NECESSARY CONDITIONS

We prove the *only if* part of Theorem 1.4. For the purpose, we prepare lemmas 4.1–4.4 whose proofs are repetitions of arguments in [10]:

LEMMA 4.1. *Let $1 \leq p, q \leq \infty$, $p < q$ and $s \in \mathbf{R}$. If $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$, then $s > n(1/p - 1/q)$.*

LEMMA 4.2. *Let $1 \leq p, q \leq \infty$ and $s \in \mathbf{R}$. If $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$, then*

$$\|\{c_k\}\|_{\ell^p} \lesssim \|\{(1 + |k|)^s c_k\}\|_{\ell^q}$$

for all finitely supported sequences $\{c_k\}_{k \in \mathbf{Z}^n}$ (that is, $c_k = 0$ except for a finite number of k 's).

PROOF OF LEMMA 4.2. Let $\eta \in \mathcal{S}(\mathbf{R}^n) \setminus \{0\}$ be such that $\text{supp } \eta \subset [-1/2, 1/2]^n$. For a finitely supported sequence $\{c_\ell\}_{\ell \in \mathbf{Z}^n}$, we set

$$f(x) = \sum_{\ell \in \mathbf{Z}^n} c_\ell e^{i\ell \cdot x} \eta(x - \ell).$$

Let $\varphi \in \mathcal{S}(\mathbf{R}^n)$ be satisfying (1). Since

$$\widehat{f}(\xi) = \sum_{\ell \in \mathbf{Z}^n} c_\ell e^{i|\ell|^2} e^{-i\ell \cdot \xi} \widehat{\eta}(\xi - \ell),$$

we see that

$$(2) \quad \varphi(D - k)f(x) = \frac{1}{(2\pi)^n} \sum_{\ell \in \mathbf{Z}^n} c_\ell e^{i|\ell|^2} \int_{\mathbf{R}^n} e^{i(x-\ell) \cdot \xi} \varphi(\xi - k) \widehat{\eta}(\xi - \ell) d\xi.$$

Using

$$\int_{\mathbf{R}^n} (1 + |x - y|)^{-M} (1 + |y|)^{-M} dy \lesssim (1 + |x|)^{-M},$$

where $M > n$, and

$$\begin{aligned} & (x - \ell)^\alpha \int_{\mathbf{R}^n} e^{i(x-\ell) \cdot \xi} \varphi(\xi - k) \widehat{\eta}(\xi - \ell) d\xi \\ &= \sum_{\alpha_1 + \alpha_2 = \alpha} C_{\alpha_1, \alpha_2} \int_{\mathbf{R}^n} e^{i(x-\ell) \cdot \xi} (\partial^{\alpha_1} \varphi)(\xi - k) (\partial^{\alpha_2} \widehat{\eta})(\xi - \ell) d\xi, \end{aligned}$$

we have

$$(3) \quad \left| \int_{\mathbf{R}^n} e^{i(x-\ell) \cdot \xi} \varphi(\xi - k) \widehat{\eta}(\xi - \ell) d\xi \right| \leq C_N (1 + |x - \ell|)^{-N} (1 + |k - \ell|)^{-N}$$

for all $N \geq 1$. Let N be a sufficiently large integer. Then, by (2) and (3),

$$|\varphi(D - k)f(x)| \lesssim \sum_{\ell \in \mathbf{Z}^n} \frac{|c_\ell|}{(1 + |x - \ell|)^N (1 + |k - \ell|)^N},$$

which provides

$$\begin{aligned} \|\varphi(D - k)f\|_{L^p} &\lesssim \sum_{\ell \in \mathbf{Z}^n} \frac{|c_\ell|}{(1 + |k - \ell|)^N} \|(1 + |\cdot - \ell|)^{-N}\|_{L^p} \\ &\approx \sum_{\ell \in \mathbf{Z}^n} \frac{|c_\ell|}{(1 + |k - \ell|)^N}. \end{aligned}$$

Then, since

$$\begin{aligned} \|f\|_{M_s^{p,q}} &= \|\{(1 + |k|)^s \|\varphi(D - k)f\|_{L^p}\}\|_{\ell^q} \\ &\lesssim \left\| \left\{ (1 + |k|)^s \sum_{\ell \in \mathbf{Z}^n} \frac{|c_\ell|}{(1 + |k - \ell|)^N} \right\} \right\|_{\ell^q} \\ &\lesssim \left\| \left\{ \sum_{\ell \in \mathbf{Z}^n} \frac{(1 + |\ell|)^s |c_\ell|}{(1 + |k - \ell|)^{N-s}} \right\} \right\|_{\ell^q}, \end{aligned}$$

we have by Young's inequality

$$(4) \quad \|f\|_{M_s^{p,q}} \lesssim \|\{(1 + |\ell|)^s c_\ell\}\|_{\ell^q}.$$

On the other hand, since $\text{supp } \eta(\cdot - \ell) \subset \ell + [-1/2, 1/2]^n$ for all $\ell \in \mathbf{R}^n$, we see that

$$\begin{aligned} (5) \quad \|f\|_{L^p} &= \left(\int_{\mathbf{R}^n} \left| \sum_{\ell \in \mathbf{Z}^n} c_\ell e^{i\ell \cdot x} \eta(x - \ell) \right|^p dx \right)^{1/p} \\ &= \left(\int_{\mathbf{R}^n} \sum_{\ell \in \mathbf{Z}^n} |c_\ell e^{i\ell \cdot x} \eta(x - \ell)|^p dx \right)^{1/p} = \|\eta\|_{L^p} \|\{c_\ell\}\|_{\ell^p} \end{aligned}$$

for $p \neq \infty$. We have easily the same conclusion for $p = \infty$. By our assumption $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$ and (4)–(5), we have

$$\|\{c_\ell\}\|_{\ell^p} \lesssim \|f\|_{L^p} \lesssim \|f\|_{M_s^{p,q}} \lesssim \|\{(1 + |\ell|)^s c_\ell\}\|_{\ell^q}.$$

The proof is complete. \square

PROOF OF LEMMA 4.1. Suppose $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$. By Lemma 4.2, we have

$$\left(\sum_{k \in \mathbf{Z}^n} |c_k|^p \right)^{1/p} \lesssim \|\{(1 + |k|)^s c_k\}\|_{\ell^q}$$

for all finitely supported sequences $\{c_k\}_{k \in \mathbf{Z}^n}$. Setting $c_k = (1+|k|)^{-s} |d_k|^{1/p}$, we see that it is equivalent to

$$\sum_{k \in \mathbf{Z}^n} (1+|k|)^{-sp} |d_k| \lesssim \|\{d_k\}\|_{\ell^{q/p}}$$

for all finitely supported sequences $\{d_k\}_{k \in \mathbf{Z}^n}$. Hence we have

$$\|\{(1+|k|)^{-sp}\}\|_{\ell^{(q/p)'}} = \sup \left| \sum_{k \in \mathbf{Z}^n} (1+|k|)^{-sp} d_k \right| \lesssim 1,$$

where the supremum is taken over all finitely supported sequences $\{d_k\}_{k \in \mathbf{Z}^n}$ such that $\|\{d_k\}\|_{\ell^{q/p}} = 1$. Note here that $(q/p)' < \infty$ from the assumption $p < q$. Hence p, q, s must satisfy $sp(q/p)' > n$, that is, $s > n(1/p - 1/q)$. \square

LEMMA 4.3. *Let $1 \leq q < p < \infty$ and $s \in \mathbf{R}$. If $L^p(\mathbf{R}^n) \hookrightarrow M_s^{p,q}(\mathbf{R}^n)$, then $s < -n(1/p + 1/q - 1)$.*

LEMMA 4.4. *Let $1 \leq p, q < \infty$ and $s \in \mathbf{R}$. If $L^p(\mathbf{R}^n) \hookrightarrow M_s^{p,q}(\mathbf{R}^n)$, then*

$$\left\{ \sum_{k \neq 0} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |\ell| \leq 2|k|} |c_\ell|^p \right)^{q/p} \right\}^{1/q} \lesssim \left(\sum_{k \neq 0} |c_k|^p \right)^{1/p}$$

for all finitely supported sequences $\{c_k\}_{k \in \mathbf{Z}^n \setminus \{0\}}$.

PROOF OF LEMMA 4.4. Let $0 < \delta < 1$ and $a \in \mathcal{S}(\mathbf{R}^n)$ be such that

$$\text{supp } a \subset [-\delta/8, \delta/8]^n, \quad \|a\|_{L^\infty} \leq 1, \quad \text{and } |\widehat{a}(\xi)| \geq C > 0 \text{ on } |\xi| \leq 2$$

(see, for example, [10, Lemma 4.3]). For a finitely supported sequence $\{c_n\}_{n \in \mathbf{Z}^n \setminus \{0\}}$, we define $f \in \mathcal{S}(\mathbf{R}^n)$ by

$$f(x) = \sum_{\ell \neq 0} c_\ell |\ell|^{n/p} a(|\ell|(x - \ell)).$$

We first estimate $\|f\|_{L^p}$. Since

$$\text{supp } a(|\ell|(\cdot - \ell)) \subset \ell + [-\delta/(8|\ell|), \delta/(8|\ell|)]^n,$$

we have

$$\begin{aligned} \|f\|_{L^p}^p &= \int_{\mathbf{R}^n} \left| \sum_{\ell \neq 0} c_\ell |\ell|^{n/p} a(|\ell|(x - \ell)) \right|^p dx \\ &= \int_{\mathbf{R}^n} \sum_{\ell \neq 0} |c_\ell|^p |\ell|^n |a(|\ell|(x - \ell))|^p dx = \|a\|_{L^p}^p \sum_{\ell \neq 0} |c_\ell|^p. \end{aligned}$$

Next, we estimate $\|f\|_{M_s^{p,q}}$. We note the following facts:

Fact 1. Let $\Psi \in \mathcal{S}(\mathbf{R}^n)$ be such that $\Psi = 1$ on $[-\delta/4, \delta/4]^n$, $\text{supp } \Psi \subset [-3\delta/8, 3\delta/8]^n$, and $|\widehat{\Psi}| \geq C > 0$ on $[-2, 2]^n$. Then we have

$$\|f\|_{M_s^{p,q}} \approx \left(\sum_{k \in \mathbf{Z}^n} (1 + |k|)^{sq} \|f * (M_k \Psi)\|_{L^p}^q \right)^{1/q},$$

where $M_k \Psi(x) = e^{ik \cdot x} \Psi(x)$.

Fact 2. For all $\ell \neq 0$, we have

$$\text{supp } a(|\ell|(\cdot - \ell)) \subset \ell + [-\delta/8|\ell|, \delta/8|\ell|]^n \subset \ell + [-\delta/8, \delta/8]^n.$$

Fact 3. For all $x \in m + [-\delta/8, \delta/8]^n$, $m \in \mathbf{Z}^n$, we have

$$\text{supp } \Psi(x - \cdot) \subset x + [-3\delta/8, 3\delta/8]^n \subset m + [-\delta/2, \delta/2]^n.$$

From these facts, we have

$$\begin{aligned} & \| (M_k \Psi) * f \|_{L^p}^p \\ & \geq \sum_{m \in \mathbf{Z}^n} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} |(M_k \Psi) * f(x)|^p dx \\ & = \sum_{m \in \mathbf{Z}^n} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} \left| \int_{\mathbf{R}^n} e^{ik \cdot (x-y)} \Psi(x-y) \sum_{\ell \neq 0} c_\ell |\ell|^{\frac{n}{p}} a(|\ell|(y-\ell)) dy \right|^p dx \\ & = \sum_{m \neq 0} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} \left| \int_{\mathbf{R}^n} e^{-ik \cdot y} \Psi(x-y) c_m |m|^{\frac{n}{p}} a(|m|(y-m)) dy \right|^p dx. \end{aligned}$$

If $x \in m + [-\delta/8, \delta/8]^n$ and $y \in \text{supp } a(|m|(\cdot - m))$, then

$$x - y \in (m + [-\delta/8, \delta/8]^n) - (m + [-\delta/8, \delta/8]^n) = [-\delta/4, \delta/4]^n,$$

and so $\Psi(x - y) = 1$. Hence,

$$\begin{aligned} & \sum_{m \neq 0} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} \left| \int_{\mathbf{R}^n} e^{-ik \cdot y} \Psi(x-y) c_m |m|^{\frac{n}{p}} a(|m|(y-m)) dy \right|^p dx \\ & = \sum_{m \neq 0} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} \left| \int_{\mathbf{R}^n} e^{-ik \cdot y} c_m |m|^{\frac{n}{p}} a(|m|(y-m)) dy \right|^p dx \\ & = \sum_{m \neq 0} \int_{m + [-\frac{\delta}{8}, \frac{\delta}{8}]^n} \left| c_m |m|^{\frac{n}{p}} |m|^{-n} \widehat{a} \left(\frac{k}{|m|} \right) \right|^p dx \\ & = \left(\frac{\delta}{4} \right)^n \sum_{m \neq 0} |c_m|^p |m|^{n-pn} \left| \widehat{a} \left(\frac{k}{|m|} \right) \right|^p. \end{aligned}$$

Moreover, using $|\widehat{a}(\xi)| \geq C > 0$ for all $1/2 \leq |\xi| \leq 2$, we obtain

$$\begin{aligned} \|(M_k \Psi) * f\|_{L^p}^p &\geq (\delta/4)^n \sum_{m \neq 0} |c_m|^p |m|^{n-pn} |\widehat{a}(k/|m|)|^p \\ &\geq (\delta/4)^n \sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p |m|^{n-pn} |\widehat{a}(k/|m|)|^p \\ &\gtrsim \sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p |m|^{n-pn} \gtrsim |k|^{n-pn} \sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p \end{aligned}$$

for all $k \neq 0$. Then

$$\begin{aligned} \|f\|_{M_s^{p,q}} &\approx \left(\sum_{k \in \mathbf{Z}^n} (1 + |k|)^{sq} \|(M_k \Psi) * f\|_{L^p}^q \right)^{1/q} \\ &\gtrsim \left\{ \sum_{k \neq 0} (1 + |k|)^{sq} \left(|k|^{n-pn} \sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p \right)^{q/p} \right\}^{1/q} \\ &\gtrsim \left\{ \sum_{k \neq 0} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p \right)^{q/p} \right\}^{1/q}. \end{aligned}$$

Therefore, by our assumption $L^p(\mathbf{R}^n) \hookrightarrow M_s^{p,q}(\mathbf{R}^n)$, we have

$$\begin{aligned} \left\{ \sum_{k \neq 0} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |m| \leq 2|k|} |c_m|^p \right)^{q/p} \right\}^{1/q} \\ \lesssim \|f\|_{M_s^{p,q}} \lesssim \|f\|_{L^p} \lesssim \left(\sum_{\ell \neq 0} |c_\ell|^p \right)^{1/p}. \end{aligned}$$

□

PROOF OF LEMMA 4.3. Suppose that $s \geq -n(1/p + 1/q - 1)$ contrary to our claim. Noting that $q/p < 1$ from the assumption $q < p$, take $\varepsilon > 0$ such that $(1 + \varepsilon)q/p < 1$ and define $\{c_k\}_{k \in \mathbf{Z}^n \setminus \{0\}}$ by

$$c_k = \begin{cases} |k|^{-n/p} (\log |k|)^{-(1+\varepsilon)/p} & \text{if } |k| \geq N, \\ 0 & \text{if } |k| < N, \end{cases}$$

where N is a sufficiently large. Note also that $\{|k|^{-n/r} (\log |k|)^{-\alpha/r}\}_{|k| \geq N} \in \ell^r$ if $\alpha > 1$, and $\{|k|^{-n/r} (\log |k|)^{-\alpha/r}\}_{|k| \geq N} \notin \ell^r$ if $\alpha \leq 1$, where $r < \infty$

(see, for example, [16, Remark 4.3]). Thus

$$\left(\sum_{k \neq 0} |c_k|^p \right)^{1/p} = \left\{ \sum_{|k| \geq N} \left(|k|^{-n/p} (\log |k|)^{-(1+\varepsilon)/p} \right)^p \right\}^{1/p} < \infty.$$

On the other hand,

$$\left\{ \sum_{k \neq 0} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |\ell| \leq 2|k|} |c_\ell|^p \right)^{q/p} \right\}^{1/q} = \infty.$$

In fact, since $n(1/p - 1) + s \geq -n/q$ and $(1 + \varepsilon)q/p < 1$, we see that

$$\begin{aligned} & \left\{ \sum_{k \neq 0} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |\ell| \leq 2|k|} |c_\ell|^p \right)^{q/p} \right\}^{1/q} \\ & \geq \left\{ \sum_{|k| \geq 2N} |k|^{(n(1/p-1)+s)q} \left(\sum_{|k|/2 \leq |\ell| \leq 2|k|} \left(|\ell|^{-n/p} (\log |\ell|)^{-(1+\varepsilon)/p} \right)^p \right)^{q/p} \right\}^{1/q} \\ & \gtrsim \left\{ \sum_{|k| \geq 2N} |k|^{(n(1/p-1)+s)q} (\log |k|)^{-(1+\varepsilon)q/p} \right\}^{1/q} \\ & \gtrsim \left\{ \sum_{|k| \geq 2N} \left(|k|^{-n/q} (\log |k|)^{-\{(1+\varepsilon)q/p\}/q} \right)^q \right\}^{1/q} = \infty. \end{aligned}$$

However, this contradicts Lemma 4.4. \square

PROOF OF THEOREM 1.4 (“ONLY IF” PART). Suppose $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$. Then we have $s \leq n\nu_2(p, q)$ by Corollary 1.2. Particularly in the case $p < q$, we have $s < -n(1/p - 1/q) = n\nu_2(p, q)$ for $p \leq 2$ by Lemma 4.1, and $s < -n(1/p' + 1/q' - 1) = n(1/p + 1/q - 1) = n\nu_2(p, q)$ for $2 \leq p$ by the dual statement of Lemma 4.3. In the case $p = \infty$, since $\mathcal{M}^{\infty,q}(\mathbf{R}^n) \hookrightarrow M^{\infty,q}(\mathbf{R}^n) \hookrightarrow L_s^\infty(\mathbf{R}^n)$, we have

$$L_{-s}^1(\mathbf{R}^n) \hookrightarrow ((L_{-s}^1(\mathbf{R}^n))^*)^* = (L_s^\infty(\mathbf{R}^n))^* \hookrightarrow (\mathcal{M}^{\infty,q}(\mathbf{R}^n))^* = M^{1,q'}(\mathbf{R}^n),$$

which means that $L^1(\mathbf{R}^n) \hookrightarrow M_s^{1,q'}(\mathbf{R}^n)$. Then we have $s < -n/q' = n(1/q - 1) = n\nu_2(\infty, q)$ for $q \neq 1$ by Lemma 2.3. All of these results yields the necessity of conditions (1)–(4). \square

5. APPLICATIONS

5.1. Unimodular Fourier multiplier. We consider the unimodular Fourier multiplier $e^{i|D|^\alpha}$, $\alpha \geq 0$, defined by

$$e^{i|D|^\alpha} f(x) = \int_{\mathbf{R}^n} e^{i|\xi|^\alpha} \widehat{f}(\xi) e^{ix \cdot \xi} d\xi, \quad f \in \mathcal{S}(\mathbf{R}^n).$$

The operator $e^{i|D|^\alpha}$ has an intimate connection with the solution $u(t, x)$ of initial value problem for the dispersive equation

$$\begin{cases} i\partial_t u + |\Delta|^{\alpha/2} u = 0, \\ u(0, x) = f(x), \end{cases}$$

$(t, x) \in \mathbf{R} \times \mathbf{R}^n$. The boundedness of $e^{i|D|^\alpha}$ on several function spaces has been studied extensively by many authors. Concerning the L^p -Sobolev spaces L_s^p and the modulation spaces $M_s^{p,q}$, the following theorems are known.

Theorem A (Miyachi [11]). *Let $1 < p < \infty$, $s \in \mathbf{R}$ and $\alpha > 1$. Then $e^{i|D|^\alpha}$ is bounded from $L_s^p(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$ if and only if $s \geq \alpha n|1/p - 1/2|$.*

Theorem B (Bényi-Gröchenig-Okoudjou-Rogers [2]). *Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $0 \leq \alpha \leq 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $M^{p,q}(\mathbf{R}^n)$.*

Theorem C (Miyachi-Nicola-Rivetti-Tabacco-Tomita [12]). *Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $\alpha > 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $M^{p,q}(\mathbf{R}^n)$ if and only if $s \geq (\alpha - 2)n|1/p - 1/2|$.*

Theorem A says that the operator $e^{i|D|^\alpha}$ is not bounded on $L^p(\mathbf{R}^n)$, and we have generally a loss of regularity of the order up to $\alpha n|1/p - 1/2|$. Theorem B describes an advantage of modulation spaces because we have no loss in the case $0 \leq \alpha \leq 2$ or smaller loss in the case $\alpha > 2$ if we consider the operator $e^{i|D|^\alpha}$ on these spaces.

Then what is the exact order of the loss when we consider the operator $e^{i|D|^\alpha}$ between L^p spaces and modulation spaces. We can answer this question by using our main theorem. The case $0 \leq \alpha \leq 2$ is rather simple, and we have the following results:

THEOREM 5.1. *Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $0 \leq \alpha \leq 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$ if and only if $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$.*

PROOF. Assume that $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$:

$$\|e^{i|D|^\alpha} f\|_{L^p} \lesssim \|f\|_{M_s^{p,q}}.$$

Note that $e^{-i|D|^\alpha}$ is also bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$. Then by taking $f = e^{-i|D|^\alpha}g$ we have

$$\|g\|_{L^p} \lesssim \|e^{-i|D|^\alpha}g\|_{M_s^{p,q}} \lesssim \|g\|_{M_s^{p,q}},$$

which means that $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$ by the equation $\overline{e^{i|D|^\alpha}f} = e^{-i|D|^\alpha}\bar{f}$. Conversely assume that $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$. Since $e^{i|D|^\alpha}$ is bounded on $M_s^{p,q}(\mathbf{R}^n)$ by Theorem B, we have

$$\|e^{i|D|^\alpha}f\|_{L^p} \lesssim \|e^{i|D|^\alpha}f\|_{M_s^{p,q}} \lesssim \|f\|_{M_s^{p,q}},$$

which means that $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$. \square

The following corollary is straightforwardly obtained from Theorem 5.1 and Theorem 1.4. The second part is just the dual statement of the first part:

COROLLARY 5.2. *Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $0 \leq \alpha \leq 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$ if and only if one of the following conditions is satisfied:*

- (1) $q \leq p < \infty$ and $s \geq -n\nu_2(p, q)$;
- (2) $p < q$ and $s > -n\nu_2(p, q)$;
- (3) $p = \infty$, $q = 1$, and $s \geq -n\nu_2(\infty, 1)$;
- (4) $p = \infty$, $q \neq 1$, and $s > -n\nu_2(\infty, q)$,

and from $L_s^p(\mathbf{R}^n)$ to $M^{p,q}(\mathbf{R}^n)$ if and only if one of the following conditions is satisfied:

- (5) $q \geq p > 1$ and $s \geq n\nu_1(p, q)$;
- (6) $p > q$ and $s > n\nu_1(p, q)$;
- (7) $p = 1$, $q = \infty$, and $s \geq n\nu_1(1, \infty)$;
- (8) $p = 1$, $q \neq \infty$ and $s > n\nu_1(1, q)$.

For $\alpha > 2$, we have the following results:

THEOREM 5.3. *Let $1 \leq p, q \leq \infty$, $s \in \mathbf{R}$ and $\alpha > 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_{s+(\alpha-2)n|1/p-1/2|}^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$ if $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$.*

PROOF. Assume that $M_s^{p,q}(\mathbf{R}^n) \hookrightarrow L^p(\mathbf{R}^n)$. Since $e^{i|D|^\alpha}$ is bounded from $M_{s+(\alpha-2)n|1/p-1/2|}^{p,q}(\mathbf{R}^n)$ to $M_s^{p,q}(\mathbf{R}^n)$ by Theorem C, we have

$$\|e^{i|D|^\alpha}f\|_{L^p} \lesssim \|e^{i|D|^\alpha}f\|_{M_s^{p,q}} \lesssim \|f\|_{M_{s+(\alpha-2)n|1/p-1/2|}^{p,q}},$$

which means that $e^{i|D|^\alpha}$ is bounded from $M_{s+(\alpha-2)n|1/p-1/2|}^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$. \square

The following corollary is obtained from Theorem 5.3, Theorem 1.4 and the duality argument again:

COROLLARY 5.4. *Let $1 \leq p, q \leq \infty, s \in \mathbf{R}$ and $\alpha > 2$. Then $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$ if one of the following conditions is satisfied:*

- (1) $q \leq p < \infty$ and $s \geq -n\nu_2(p, q) + (\alpha - 2)n|1/p - 1/2|$;
- (2) $p < q$ and $s > -n\nu_2(p, q) + (\alpha - 2)n|1/p - 1/2|$;
- (3) $p = \infty, q = 1$, and $s \geq -n\nu_2(\infty, 1) + (\alpha - 2)n|1/p - 1/2|$;
- (4) $p = \infty, q \neq 1$, and $s > -n\nu_2(\infty, q) + (\alpha - 2)n|1/p - 1/2|$,

and from $L_s^p(\mathbf{R}^n)$ to $M^{p,q}(\mathbf{R}^n)$ if one of the following conditions is satisfied:

- (5) $q \geq p > 1$ and $s \geq n\nu_1(p, q) + (\alpha - 2)n|1/p - 1/2|$;
- (6) $p > q$ and $s > n\nu_1(p, q) + (\alpha - 2)n|1/p - 1/2|$;
- (7) $p = 1, q = \infty$, and $s \geq n\nu_1(1, \infty) + (\alpha - 2)n|1/p - 1/2|$;
- (8) $p = 1, q \neq \infty$ and $s > n\nu_1(1, q) + (\alpha - 2)n|1/p - 1/2|$.

Although the converse of Theorem 5.3 is not true, we believe that the converse of Corollary 5.4 is still true. In fact we have at least the following result:

THEOREM 5.5. *Let $1 \leq p, q \leq \infty, s \in \mathbf{R}$ and $\alpha > 2$. Suppose that $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$. Then we have $s \geq -n\nu_2(p, q) + (\alpha - 2)n|1/p - 1/2|$. Suppose that $e^{i|D|^\alpha}$ is bounded from $L_s^p(\mathbf{R}^n)$ to $M^{p,q}(\mathbf{R}^n)$ instead. Then we have $s \geq n\nu_1(p, q) + (\alpha - 2)n|1/p - 1/2|$.*

To prove Theorem 5.5, we use the following lemma.

LEMMA 5.6. *Let $1 \leq p, q \leq \infty, s \in \mathbf{R}$ and $\alpha \geq 0$. Then*

$$(6) \quad \sup_{k \in \mathbf{Z}^n} \langle k \rangle^{-s} \|\varphi(D - k)e^{i|D|^\alpha}\|_{L^p \rightarrow L^p} \lesssim \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p},$$

where φ is a function satisfying (1).

PROOF. Let N be a positive integer such that

$$\varphi(\cdot - k) = \sum_{|\ell| \leq N} \varphi(\cdot - k)\varphi(\cdot - (k + \ell))$$

for all $k \in \mathbf{Z}^n$. Then we have

$$\begin{aligned}
& \|\varphi(D - k)e^{i|D|^\alpha}f\|_{L^p} \\
& \leq \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p} \|\varphi(D - k)f\|_{M_s^{p,q}} \\
& = \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p} \left(\sum_{m \in \mathbf{Z}^n} \langle m \rangle^{sq} \|\varphi(D - m)\varphi(D - k)f\|_{L^p}^q \right)^{1/q} \\
& = \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p} \left(\sum_{|\ell| \leq N} \langle k + \ell \rangle^{sq} \|\varphi(D - (k + \ell))\varphi(D - k)f\|_{L^p}^q \right)^{1/q} \\
& \lesssim \langle k \rangle^s \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p} \|f\|_{L^p}
\end{aligned}$$

for $f \in \mathcal{S}(\mathbf{R}^n)$ and $q \neq \infty$. We have easily the same conclusion for $q = \infty$. Hence, we obtain the desired result. \square

REMARK 5.7. We remark that since

$$\|e^{i|D|^\alpha}\|_{M_s^{p,\tilde{q}} \rightarrow M^{p,\tilde{q}}} \approx \sup_{k \in \mathbf{Z}^n} \langle k \rangle^{-s} \|\varphi(D - k)e^{i|D|^\alpha}\|_{L^p \rightarrow L^p}$$

for $1 \leq p, \tilde{q} \leq \infty, s \in \mathbf{R}$ (see [12, Lemma 2.2]), we have

$$\|e^{i|D|^\alpha}\|_{M_s^{p,\tilde{q}} \rightarrow M^{p,\tilde{q}}} \lesssim \|e^{i|D|^\alpha}\|_{M_s^{p,q} \rightarrow L^p}$$

for all $1 \leq p, q, \tilde{q} \leq \infty$.

Now, we prove Theorem 5.5.

Proof of Theorem 5.5. Since the latter is just the dual statement of the former, we prove only the former. Suppose that $e^{i|D|^\alpha}$ is bounded from $M_s^{p,q}(\mathbf{R}^n)$ to $L^p(\mathbf{R}^n)$:

$$(7) \quad \|e^{i|D|^\alpha}f\|_{L^p} \lesssim \|f\|_{M_s^{p,q}}, \quad f \in \mathcal{S}(\mathbf{R}^n).$$

(i) Let $q \leq \min(p, p')$. By the necessary condition of Theorem C and Remark 5.7, we have $s \geq (\alpha - 2)n|1/p - 1/2|$. Since $\nu_2(p, q) = 0$, we obtain the desired result.

(ii) Let $1 \leq p \leq 2$ and $p \leq q \leq p'$. Note that inequality (7) can be written as

$$(8) \quad \|e^{i|D|^\alpha} \langle D \rangle^{-s} f\|_{L^p} \lesssim \|f\|_{M^{p,q}}, \quad f \in \mathcal{S}(\mathbf{R}^n)$$

by the lifting property. Here, we denote $\langle D \rangle^{-s} f = (\langle \cdot \rangle^{-s} \widehat{f}(\cdot))^\vee$ for $s \in \mathbf{R}$.

Let $g \in \mathcal{S}(\mathbf{R}^n)$ be such that

$$(9) \quad \text{supp } \widehat{g} \subset \{\xi \mid 2^{-1} < |\xi| < 2\} \text{ and } \widehat{g}(\xi) = 1 \text{ on } \{\xi \mid 2^{-1/2} < |\xi| < 2^{1/2}\},$$

and test (8) with a specific $f = U_\lambda g$, $\lambda \geq 1$. Since

$$e^{i|D|^\alpha} \langle D \rangle^{-s} U_\lambda g = U_\lambda (e^{i|\lambda D|^\alpha} \langle \lambda D \rangle^{-s} g),$$

it follows from Theorem 2.2 that

$$\lambda^{-n/p} \|e^{i|\lambda D|^\alpha} \langle \lambda D \rangle^{-s} g\|_{L^p} \lesssim \lambda^{n\mu_1(p,q)} \|g\|_{M^{p,q}}.$$

On the other hand, by the change of variable $x \mapsto \lambda^\alpha x$ and the method of stationary phase, we obtain

$$\begin{aligned} \|e^{i|\lambda D|^\alpha} \langle \lambda D \rangle^{-s} g\|_{L^p} &= \left\| \int_{\mathbf{R}^n} e^{ix \cdot \xi + i|\lambda \xi|^\alpha} \langle \lambda \xi \rangle^{-s} \widehat{g}(\xi) d\xi \right\|_{L^p} \\ &= \lambda^{\alpha n/p} \left\| \int_{\mathbf{R}^n} e^{i\lambda^\alpha (x \cdot \xi + |\xi|^\alpha)} \langle \lambda \xi \rangle^{-s} \widehat{g}(\xi) d\xi \right\|_{L^p} \\ &\gtrsim \lambda^{\alpha n/p - \alpha n/2 - s}. \end{aligned}$$

Combining these two estimates, we get

$$\lambda^{n\mu_1(p,q) + n/p - \alpha n/2 + s} \gtrsim 1$$

for all $\lambda \geq 1$. Letting $\lambda \rightarrow \infty$ yields the necessary condition

$$\begin{aligned} s &\geq -n(1/q - 1) - n/p + \alpha n/p - \alpha n/2 \\ &= (\alpha - 2)n(1/p - 1/2) + n/p - n/q \\ &= (\alpha - 2)n|1/p - 1/2| - n\nu_2(p, q), \end{aligned}$$

since $\mu_1(p, q) = 1/q - 1$ and $\nu_2(p, q) = -1/p + 1/q$.

(iii) Let $2 \leq p \leq \infty$ and $p' \leq q \leq p$. By duality, we have

$$\|e^{-i|D|^\alpha} f\|_{L^{p'}} \lesssim \|f\|_{M_{-s}^{p',q'}}, \quad f \in \mathcal{S}(\mathbf{R}^n).$$

So, we have only to prove the following lemma.

LEMMA 5.8. *Let $1 \leq p' \leq 2, p' \leq q' \leq p$ and $s \in \mathbf{R}$. If $e^{i|D|^\alpha}$ is bounded from $L_s^{p'}(\mathbf{R}^n)$ to $M^{p',q'}(\mathbf{R}^n)$, then $s \geq (\alpha - 2)n|1/p - 1/2| - n\nu_2(p, q)$.*

Proof of Lemma 5.8. Set $f = U_\lambda g$, $\lambda \geq 1$, where g is a function satisfying (9). Then, by Lemma 2.2, we have

$$\begin{aligned} \|e^{i|D|^\alpha} f\|_{M^{p',q'}} &= \|e^{i|D|^\alpha} U_\lambda g\|_{M^{p',q'}} \\ &= \|U_\lambda (e^{i|\lambda D|^\alpha} g)\|_{M^{p',q'}} \\ &\gtrsim \lambda^{n\mu_2(p',q')} \|e^{i|\lambda D|^\alpha} g\|_{M^{p',q'}}. \end{aligned}$$

In the same way as (ii), we obtain, by the change of variable $x \mapsto \lambda^\alpha x$ and the method of stationary phase,

$$\begin{aligned}
\|e^{i|\lambda D|^\alpha} g\|_{M^{p',q'}} &= \left(\sum_{k \in \mathbf{Z}^n} \|\varphi(D-k)e^{i|\lambda D|^\alpha} g\|_{L^{p'}}^{q'} \right)^{1/q'} \\
&\gtrsim \left\| \int_{\mathbf{R}^n} \varphi(\xi) e^{ix \cdot \xi + i|\lambda \xi|^\alpha} \widehat{g}(\xi) d\xi \right\|_{L^{p'}} \\
&= \lambda^{\alpha n/p'} \left\| \int_{\mathbf{R}^n} e^{i\lambda^\alpha (x \cdot \xi + |\xi|^\alpha)} \varphi(\xi) \widehat{g}(\xi) d\xi \right\|_{L^{p'}} \\
&\gtrsim \lambda^{\alpha n/p' - \alpha n/2}.
\end{aligned}$$

Hence, we have

$$\|e^{i|D|^\alpha} f\|_{M^{p',q'}} \gtrsim \lambda^{n\mu_2(p',q')} \lambda^{\alpha n(1/p' - 1/2)}.$$

On the other hand, we have

$$\|f\|_{L_s^{p'}} = \|U_\lambda g\|_{L_s^{p'}} \approx \lambda^{-n/p'} \lambda^s.$$

Combining these two estimates, we obtain


$$\lambda^{s - n/p' - n\mu_2(p',q') - \alpha n(1/p' - 1/2)} \gtrsim 1$$

for all $\lambda \geq 1$. Letting $\lambda \rightarrow \infty$ yields the necessary condition

$$\begin{aligned}
s &\geq \alpha n(1/p' - 1/2) + n/p' + n(-2/p' + 1/q') \\
&= (\alpha - 2)n(1/p' - 1/2) + 2n(1/p' - 1/2) + n/p' + n(-2/p' + 1/q') \\
&= (\alpha - 2)n(1/p' - 1/2) + n(1/p' + 1/q' - 1) \\
&= (\alpha - 2)n(1/2 - 1/p) + n(1 - 1/p - 1/q) \\
&= (\alpha - 2)n|1/p - 1/2| - n\nu_2(p, q),
\end{aligned}$$

since $\mu_2(p', q') = -2/p' + 1/q'$ and $\nu_2(p, q) = 1/p + 1/q - 1$. \square

(iv) Let $2 \leq p \leq \infty$ and $p < q$. Contrary to our claim, suppose that there exists $\varepsilon > 0$ such that $s = (\alpha - 2)n|1/p - 1/2| - n\nu_2(p, q) - \varepsilon$ implies (7). Then, by interpolation with the estimate for a point $Q(1/p_1, 1/q_1)$ with $2 < p_1 < \infty$, $p'_1 < q_1 < p_1$ and $s = (\alpha - 2)n|1/p_1 - 1/2| - n\nu_2(p_1, q_1)$ (which holds by Corollary 5.4), one would obtain an improved estimates of the segment joining $P(1/p, 1/q)$ and $Q(1/p_1, 1/q_1)$, which is not possible. In the same way as above, we can treat the case $1 \leq p \leq 2$ and $p' < q$, and we have the conclusion.



□

5.2. Pseudo-differential operator. Let $\sigma(x, \xi)$ be a function on $\mathbf{R}^n \times \mathbf{R}^n$. Then the pseudo-differential operator $\sigma^W(X, D)$ is defined by

$$\sigma^W(X, D)f(x) = \frac{1}{(2\pi)^n} \iint_{\mathbf{R}^{2n}} \sigma\left(\frac{x+y}{2}, \xi\right) e^{i(x-y)\cdot\xi} f(y) dy d\xi.$$

We give a condition for $\sigma^W(X, D)$ to be bounded from $M^{p,q}(\mathbf{R}^n)$ to $L_s^p(\mathbf{R}^n)$. To this end, we recall the following fundamental results about $\sigma^W(X, D)$ with $\sigma \in M^{\infty,1}(\mathbf{R}^{2n})$.

Theorem D (Sjöstrand [13],[14]). *Let $\sigma \in M^{\infty,1}(\mathbf{R}^{2n})$. Then $\sigma^W(X, D)$ is bounded on $L^2(\mathbf{R}^n)$. Moreover, if $\sigma^W(X, D)$ is invertible on $L^2(\mathbf{R}^n)$, then $(\sigma^W(X, D))^{-1} = \tau^W(X, D)$ for some $\tau \in M^{\infty,1}(\mathbf{R}^{2n})$.*

Theorem E (Gröchenig-Heil [8]). *Let $1 \leq p, q \leq \infty$ and $\sigma \in M^{\infty,1}(\mathbf{R}^{2n})$. Then $\sigma^W(X, D)$ is bounded on $M^{p,q}(\mathbf{R}^n)$.*

Now, we state our result which is an analog of Theorems 5.1 and 5.3.

THEOREM 5.9. *Let $1 \leq p, q \leq \infty, s \in \mathbf{R}^n$ and $\sigma \in M^{\infty,1}(\mathbf{R}^{2n})$.*

- (1) *If $M^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$, then $\sigma^W(X, D)$ is bounded from $M^{p,q}(\mathbf{R}^n)$ to $L_s^p(\mathbf{R}^n)$.*
- (2) *Suppose that $\sigma^W(X, D)$ is invertible on $L^2(\mathbf{R}^n)$. If $\sigma^W(X, D)$ is bounded from $M^{p,q}(\mathbf{R}^n)$ to $L_s^p(\mathbf{R}^n)$, then $\mathcal{M}^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$.*

PROOF. (1) Since $\sigma^W(X, D)$ is bounded on $M^{p,q}(\mathbf{R}^n)$ by Theorem E, we have

$$\|\sigma^W(X, D)f\|_{L_s^p} \lesssim \|\sigma^W(X, D)f\|_{M^{p,q}} \lesssim \|f\|_{M^{p,q}},$$

which means $\sigma^W(X, D)$ is bounded from $M^{p,q}(\mathbf{R}^n)$ to $L_s^p(\mathbf{R}^n)$.

(2) Since $(\sigma^W(X, D))^{-1} = \tau^W(X, D)$ for some $\tau \in M^{\infty,1}(\mathbf{R}^{2n})$ and $\tau^W(X, D)$ is bounded on $M^{p,q}(\mathbf{R}^n)$ by Theorems D and E, we obtain

$$\|f\|_{L_s^p} = \|\sigma^W(X, D)\tau^W(X, D)f\|_{L_s^p} \lesssim \|\tau^W(X, D)f\|_{M^{p,q}} \lesssim \|f\|_{M^{p,q}}$$

for $f \in \mathcal{S}(\mathbf{R}^n) \subset L^2(\mathbf{R}^n)$, which means $\mathcal{M}^{p,q}(\mathbf{R}^n) \hookrightarrow L_s^p(\mathbf{R}^n)$. \square

If we combine Theorem 5.9 and Theorem 1.4, we have similar results to Corollaries 5.2 and 5.4. We will not state them explicitly but it is just a straightforward task.

REFERENCES

- [1] Á. Bényi, K. Gröchenig, C. Heil and K. Okoudjou, *Modulation spaces and a class of bounded multilinear pseudodifferential operators*, J. Operator Theory 54 (2005), pp. 387–399.
- [2] Á. Bényi, K. Gröchenig, K. Okoudjou and L.G. Rogers, *Unimodular Fourier multipliers for modulation spaces*, J. Funct. Anal. 246 (2007), pp. 366–384.
- [3] E. Cordero, K. Gröchenig, *Time-frequency analysis of localization operators*, J. Funct. Anal. 205 (2003), pp. 107–131.
- [4] H.G. Feichtinger, *Modulation spaces on locally compact abelian groups*, in: M. Krishna, R. Radha and S. Thangavelu (Eds.), Wavelets and their Applications, Chennai, India, Allied Publishers, New Delhi, 2003, pp. 99–140, Updated version of a technical report, University of Vienna, 1983.
- [5] D. Goldberg, *A local version of real Hardy spaces*, Duke Math. J. 46 (1979), pp. 27–42.
- [6] K. Gröchenig, *Foundations of Time-Frequency Analysis*, Birkhäuser, 2001.
- [7] K. Gröchenig, *Time-frequency analysis of Sjöstrand’s class*, Rev. Mat. Iberoam. 22 (2006), pp. 703–724.
- [8] K. Gröchenig and C. Heil, *Modulation spaces and pseudodifferential operators*, Integral Equations Operator Theory 34 (1999), pp. 439–457.
- [9] C. Heil, J. Ramanathan and P. Topiwala, *Singular values of compact pseudodifferential operators*, J. Funct. Anal. 150 (1997), pp. 426–452.
- [10] M. Kobayashi, A. Miyachi and N. Tomita, *Embedding relations between local Hardy and modulation spaces*, Studia Math. 192 (2009), pp. 79–96.
- [11] A. Miyachi, *On some Fourier multipliers for $H^p(\mathbf{R}^n)$* , J. Fac. Sci. Univ. Tokyo Sect. IA Math. 27 (1980), pp. 157–179.
- [12] A. Miyachi, F. Nicola, S. Rivetti, A. Tabacco and N. Tomita, *Estimates for unimodular Fourier multipliers on modulation spaces*, Proc. Amer. Math. Soc. 137 (2009), pp. 3869–3883.
- [13] J. Sjöstrand, *An algebra of pseudodifferential operators*, Math. Res. Lett. 1 (1994), pp. 185–192.
- [14] J. Sjöstrand, *Wiener type algebras of pseudodifferential operators*, Séminaire sur les Équations aux Dérivées Partielles, 1994-1995, Exp.No. IV, 21 pp., École Polytech., Palaiseau, 1995.
- [15] M. Sugimoto and N. Tomita, *The dilation property of modulation spaces and their inclusion relation with Besov spaces*, J. Funct. Anal. 248 (2007), pp. 79–106.
- [16] M. Sugimoto and N. Tomita, *A remark on fractional integrals on modulation spaces*, Math. Nachr. 281 (2008), pp. 1372–1379.
- [17] K. Tachizawa, *The boundedness of pseudodifferential operators on modulation spaces*, Math. Nachr. 168 (1994), pp. 263–277.

- [18] J. Toft, *Continuity properties for modulation spaces, with applications to pseudo-differential calculus, I*, J. Funct. Anal. 207 (2004), pp. 399–429.
- [19] J. Toft, *Convolutions and embeddings for weighted modulation spaces*, Advances in pseudo-differential operators, 165–186, Oper. Theory Adv. Appl, 155, Birkhäuser, 2004.
- [20] H. Triebel, *Theory of function spaces. II*, Birkhäuser, 1992.
- [21] B. Wang and C. Huang, *Frequency-uniform decomposition method for the generalized BO, KdV and NLS equations*, J. Differential Equations 239 (2007), pp. 213–250.
- [22] B. Wang and H. Hudzik, *The global Cauchy problem for the NLS and NLKG with small rough data*, J. Differential Equations 232 (2007), pp. 36–73.
- [23] B. Wang, L. Zhao and B. Guo, *Isometric decomposition operators, function spaces $E_{p,q}^\lambda$ and applications to nonlinear evolution equations*, J. Funct. Anal. 233 (2006), pp. 1–39.

MASAHARU KOBAYASHI, DEPARTMENT OF MATHEMATICS, TOKYO UNIVERSITY OF SCIENCE, KAGURAZAKA 1-3, SHINJUKU-KU, TOKYO 162-8601, JAPAN
E-mail address: kobayashi@jan.rikadai.jp

MITSURU SUGIMOTO, GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY,, FUROCHO, CHIKUSA-KU, NAGOYA 464-8602, JAPAN
E-mail address: sugimoto@math.nagoya-u.ac.jp